

NAME: _____ ()

CLASS: _____



FAIRFIELD METHODIST SCHOOL (SECONDARY)
PRELIMINARY EXAMINATION 2024
SECONDARY 4 EXPRESS

ADDITIONAL MATHEMATICS**4049/01****Paper 1****Date: 21 August 2024****Duration: 2 hours 15 minutes**

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your name, index number and class on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Answer **all** the questions.

The number of marks is given in brackets [] at the end of each question or part question.

If working is needed for any question it must be shown with the answer.

Omission of essential working will result in loss of marks.

The total of the marks for this paper is 90.

The use of an approved scientific calculator is expected, where appropriate.

If the degree of accuracy is not specified in the question and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.

For π , use either your calculator value or 3.142.**For Examiner's Use**

Table of Penalties		Question Number	Parent's/Guardian's Signature	90
Presentation	☐ 1 ☐ 2			
Rounding off	☐ 1			

This question paper consists of **22** printed pages

NAME: _____ ()

CLASS: _____

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial expansion

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\dots(n-r+1)}{r!}$

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$



NAME: _____ ()

CLASS: _____

Answer **all** the questions.

- 1 (i) Express $4x^2 + 8x - 5$ in the form $p(x+q)^2 + r$, where p , q and r are constants to be found. [3]

- (ii) Hence, state the coordinates of the turning point of the curve $y = 4x^2 + 8x - 5$. [1]

NAME: _____ ()


CLASS: _____

- 2 Without using a calculator, find the values of a and b for which the solution of the equation $x\sqrt{24} = x\sqrt{3} + \sqrt{6}$ is $\frac{a+\sqrt{b}}{7}$. [5]

NAME: _____ ()

CLASS: _____

- 3 Points A , B , C and D lie on a circle. The tangent to the circle at A meets BD produced at E . AE is parallel to BC .



Prove that

(i) $AB = AC$,

[3]


(ii) $\angle CDE = 2\angle ABC$.

[3]

NAME: _____ ()

CLASS: _____

- 4 (a) The diagram shows part of the graph of $y = a \tan bx + c$. The graph has vertical asymptotes at $x = -4\pi$ and $x = 4\pi$ and passes through the points P and Q .



- (i) Explain why $b = \frac{1}{8}$. [2]

- (ii) Hence find the equation of the curve. [2]



NAME: _____ ()

CLASS: _____

- (b) The function $f(x)$ is defined by $f(x) = 4 + 3\sin 2x$ for $0^\circ \leq x \leq 360^\circ$.
Sketch the graph of $y = f(x)$ on the axes below.


[2]



NAME: _____ ()

CLASS: _____

- 5 The diagram shows a shape made by cutting an equilateral triangle out of a rectangle of width x cm.



The perimeter of the shape is 20 cm.

- (i) Show that the area, A cm², of the shape is given by $A = 10x - \left(\frac{6 + \sqrt{3}}{4}\right)x^2$. [3]

NAME: _____ ()

CLASS: _____

- (ii) Given that x can vary, find the value of x which produces the maximum area and calculate this maximum area. Give your answers to 2 significant figures. [4]

NAME: _____ ()

CLASS: _____

6 (a) Express $\frac{8x+13}{(1+2x)(2+x)^2}$ in partial fractions.

[5]

NAME: _____ ()

CLASS: _____

(b) Hence, evaluate $\int_1^2 \frac{8x+13}{(1+2x)(2+x)^2} dx$.

[3]

NAME: _____ ()

CLASS: _____

- 7 The polynomial $f(x)$ is such that $f(x) = 6x^3 + ax^2 - 50x + b$, where a and b are integers. It is given that $f(x)$ is divisible by $2x - 3$ and that $f'(1) = 6$.

(a) Find the values of a and b .

[5]



NAME: _____ ()

CLASS: _____

(b) Using your values of a and b , solve the equation $f(x) = 0$.

[3]

NAME: _____ ()

CLASS: _____

- 8 (a) Show that the solution of the equation $2^{3x+4} \times 5^{2x-1} = 16^x \times 5^{3x}$ is $\lg \frac{16}{5}$. [3]

- (b) Express $2\log_2 x - \log_2(x-4) = 3$ as a quadratic equation in x and explain why there are no real solutions. [5]



NAME: _____ ()

CLASS: _____

- 9 (a) The variables x and y increase in such a way that when $x = 3$, the rate of increase of y with respect to time is three times the rate of increase of x with respect to time.
Given that $y = k\sqrt{3x+7}$, where k is a constant, find the value of k . [4]

NAME: _____ ()

CLASS: _____

- (b) The mass, m grams, of a radioactive sample, present at time t days after being observed, is given by $m = 24e^{-0.02t}$.

Find

- (i) the initial mass of the radioactive sample,

[1]

- (ii) the time taken for the sample to decrease to half its initial mass,

[2]

- (iii) the rate at which the mass is decreasing after 12 hours.

[2]

NAME: _____ ()

CLASS: _____

10 (i) Show that $\frac{\tan x}{1 + \sec x} + \frac{1 + \sec x}{\tan x} = \frac{2}{\sin x}$. [5]

NAME: _____ ()


CLASS: _____

(ii) Hence solve the equation $\frac{\tan x}{1+\sec x} + \frac{1+\sec x}{\tan x} = 1+3\sin x$ for $0^\circ \leq x \leq 360^\circ$. [4]

NAME: _____ ()

CLASS: _____

- 11 The diagram shows part of the curve $y = (9-x)(x-3)$ and the line $y = k-3$, where $k > 3$. The line through the maximum point of the curve, parallel to the y -axis, meets the x -axis at A . The curve meets the x -axis at B and the line $y = k-3$ meets the curve at the point $C(k, k-3)$.



- (i) Show that the value of k is 8.

[4]

NAME: _____ ()

CLASS: _____

(ii) Find the area of the shaded region.


[5]

NAME: _____ ()

CLASS: _____

- 12 Solutions to this question by accurate drawing will not be accepted.

The diagram (not drawn to scale) shows a trapezium $OPQR$ in which PQ is parallel to OR and $\angle ORQ = 90^\circ$. The coordinates of P and R are $(-4, 3)$ and $(4, 2)$ respectively and O is the origin.



- (i) Find the coordinates of Q .

[5]

NAME: _____ ()

CLASS: _____

- (ii) PQ meets the y -axis at T . Show that triangle ORT is isosceles.

[3]

The point S is such that $ORPS$ forms a parallelogram.

- (iii) Find the coordinates of S .

[1]

- (iv) Find the area of the trapezium $OPQR$.

[2]

~ End of Paper ~

FMS(S) SEC 4Exp Additional Mathematics Paper 1 Preliminary Exam 2024 Marking Scheme

1i	$4x^2 + 8x - 5 = 4(x^2 + 2x) - 5$ $= 4[(x+1)^2 - 1] - 5$ $= 4(x+1)^2 - 9$	M1 M1 complete square A1 B1	AO1
ii	Turning point = $(-1, -9)$	B1	AO1

2	$x\sqrt{24} = x\sqrt{3} + \sqrt{6}$ $x(\sqrt{24} - \sqrt{3}) = \sqrt{6}$ $x = \frac{\sqrt{6}}{(\sqrt{24} - \sqrt{3})}$ $x = \frac{\sqrt{6}}{(2\sqrt{6} - \sqrt{3})} \times \frac{2\sqrt{6} + \sqrt{3}}{2\sqrt{6} + \sqrt{3}}$ $= \frac{2(6) + \sqrt{18}}{4(6) - 3}$ $= \frac{12 + 3\sqrt{2}}{21}$ $= \frac{4 + \sqrt{2}}{7}$ $a = 4 \text{ and } b = 2$	M1 factorise x M1 Multiply by conjugate surd M1 simplification A1, A1	AO1
---	---	--	-----

3i	$\angle CAE = \angle ABC$ (alt segment theorem/tangent chord thm) $\angle ACB = \angle CAE$ (alternate $\angle$ s, $BC \parallel AE$) $\angle ABC = \angle ACB$ (base $\angle$ s of isos Δ) $\therefore AB = AC$ (shown)	B1 B1 AG1	AO3
3ii	$\angle BAC = 180^\circ - 2\angle ABC$ ($\angle$ sum of Δ) $\angle BDC = \angle BAC$ ($\angle$ s in same segment) $\angle CDE = 180^\circ - \angle BDC$ (adj $\angle$ s on a straight line) $\angle CDE = 180^\circ - (180^\circ - 2\angle ABC)$ (adj $\angle$ s on a straight line) $\angle CDE = 2\angle ABC$ (shown)	M1 B1 AG1	AO3

4a i	Since the period = 8π $8\pi = \frac{\pi}{b}$ $b = \frac{1}{8}$	B1 AG1	AO3
ii	$c = 3$ $7 = a \tan\left(\frac{\pi}{4}\right) + 3$ $a = 4$ $y = 4 \tan\left(\frac{x}{8}\right) + 3$	M1 A1	AO1
4b		B1 correct sinusoidal shape with correct turning points B1 two cycles	AO1

5i	Length of rectangle = $\frac{20-3x}{2}$ Area = $\left(\frac{20-3x}{2}\right)x - \frac{1}{2}x^2 \sin 60^\circ$ $= 10x - \frac{3}{2}x^2 - \frac{1}{2}x^2 \frac{\sqrt{3}}{2}$ $= 10x - \left(\frac{6+\sqrt{3}}{4}\right)x^2$	B1 M1 AG1	AO3
5ii	$\frac{dA}{dx} = 10 - 2\left(\frac{6+\sqrt{3}}{4}\right)x$ For stationary point, $10 - 2\left(\frac{6+\sqrt{3}}{4}\right)x = 0$ $(6+\sqrt{3})x = 20$ $x = \frac{20}{(6+\sqrt{3})} = 2.5866 = 2.6 \text{ (2 s.f.)}$ $A = 10(2.5866) - \left(\frac{6+\sqrt{3}}{4}\right)(2.5866)^2 = 12.933$	B1 - diff M1 - equate to 0 A1 A1	AO1

$A = 13$		
6a	Let $\frac{8x+13}{(1+2x)(2+x)^2} = \frac{A}{1+2x} + \frac{B}{2+x} + \frac{C}{(2+x)^2}$ $8x+13 = A(2+x)^2 + B(1+2x)(2+x) + C(1+2x)$ $x = -2 \Rightarrow -3 = -3C$ $C = 1$ $x = -\frac{1}{2} \Rightarrow 9 = \frac{9}{4}A$ $A = 4$ $x = 0 \Rightarrow 13 = 4A + 2B + C$ $13 = 4(4) + 2B + 1$ $B = -2$ $\therefore \frac{8x+13}{(1+2x)(2+x)^2} = \frac{4}{(1+2x)} - \frac{2}{(2+x)} + \frac{1}{(2+x)^2}$	AO1 B1 M1 – sub or compare coefficient A1 (anyone of A, B or C correct) A1 (all 3 values A, B & C) A1
6b	$\int_1^2 \frac{8x+13}{(1+2x)(2+x)^2} dx$ $= \int_1^2 \left[\frac{4}{(1+2x)} - \frac{2}{(2+x)} + \frac{1}{(2+x)^2} \right] dx$ $= \left[\frac{4}{2} \ln(1+2x) - 2 \ln(2+x) - \frac{1}{2+x} \right]_1^2$ $= \left[2 \ln 5 - 2 \ln 4 - \frac{1}{4} \right] - \left[2 \ln 3 - 2 \ln 3 - \frac{1}{3} \right]$ $= 2 \ln \frac{5}{4} + \frac{1}{12}$ or $\ln \frac{25}{16} + \frac{1}{12}$ or 0.530 (to 3 sig fig)	AO2 B2 [B1 for 2 correct ln term; B1 for the 3rd term] B1

7a	$f'(x) = 18x^2 + 2ax - 50$ Since $f'(1) = 6$ $18 + 2a - 50 = 6$ $a = 19$ Given $f\left(\frac{3}{2}\right) = 0$ $6\left(\frac{3}{2}\right)^3 + 19\left(\frac{3}{2}\right)^2 - 50\left(\frac{3}{2}\right) + b = 0$ $20.25 + 42.75 - 75 + b = 0$	B1 M1 forming equation A1 M1 forming equation A1	AO2
----	---	---	-----

[illegible]

8a	$2^{3x+4} \times 5^{2x-1} = 16^x \times 5^{3x}$ $2^{3x} \times 2^4 \times 5^{2x} \times 5^{-1} = 2^{4x} \times 5^{3x}$ $\frac{16}{5} = 2^{4x} \div 2^{3x} \times 5^{3x} \div 5^{2x}$ $\frac{16}{5} = 2^x \times 5^x$ $\frac{16}{5} = 10^x$ $\lg \frac{16}{5} = \lg 10^x$ $x = \lg \frac{16}{5} \text{ (shown)}$	M1: simplify base 2 and base 5 terms correctly M1: introduce log on both sides AG1	AO3
8b	$2 \log_2 x - \log_2 (x-4) = 3$ $\log_2 x^2 - \log_2 (x-4) = 3$ $\log_2 \frac{x^2}{x-4} = 3$ $\frac{x^2}{x-4} = 2^3$ $x^2 = 8(x-4)$ $x^2 - 8x + 32 = 0$ $x = \frac{-(-8) \pm \sqrt{(-8)^2 - 4(1)(32)}}{2(1)}$	M1: power/quotient law M1: change to index form M1: form quadratic eqn M1 solve for x OR show $b^2 - 4ac = -64$	AO3

$x = \frac{8 \pm \sqrt{-64}}{2}$ $\therefore$ No real solution (shown)	AG1: explain $\sqrt{-64}$ does not exist or $b^2 - 4ac < 0 \therefore$ no real solution	
---	--	--


9a	$y = k\sqrt{3x+7},$ $\frac{dy}{dx} = \frac{3k}{2}(3x+7)^{-\frac{1}{2}}$ Given $\frac{dy}{dt} = 3 \frac{dx}{dt}$ $\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$ $\frac{3dx}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$ $3 = \frac{3k}{2}(3x+7)^{-\frac{1}{2}}$ Sub $x = 3$ $1 = \frac{k}{2}(3 \times 3 + 7)^{-\frac{1}{2}}$ $1 = \frac{k}{2} \times \frac{1}{4}$ $k = 8$	B1: correct differentiation M1: write the correct ratio M1: form chain rule correctly and sub $x = 3$ A1	AO2
9b i	Sub $t = 0, m = 24 \text{ g}$	B1	AO1
ii	$24e^{-0.02t} = 12$ $\ln e^{-0.02t} = \ln 0.5$ $-0.02t = \ln 0.5$ $t = \frac{\ln 0.5}{-0.02} = 34.7 \text{ days}$	M1 A1	AO1
iii	$\frac{dm}{dt} = -0.48e^{-0.02t}$ Sub $t = 0.5, \frac{dm}{dt} = -0.48e^{-0.02(0.5)} = -0.475 \text{ (3 s.f.)}$ Mass is decreasing at 0.475 g/day	B1 B1 (must write statement)	AO1

10 i	LHS $= \left[\frac{\sin x}{\cos x} \div \left(1 + \frac{1}{\cos x} \right) \right] + \left[\left(1 + \frac{1}{\cos x} \right) \div \frac{\sin x}{\cos x} \right]$ $= \left[\frac{\sin x}{\cos x} \times \left(\frac{\cos x}{\cos x + 1} \right) \right] + \left[\left(\frac{\cos x + 1}{\cos x} \right) \times \frac{\cos x}{\sin x} \right]$ $= \frac{\sin x}{\cos x + 1} + \frac{1 + \cos x}{\sin x}$ $= \frac{\sin^2 x + (1 + \cos x)^2}{\sin x(1 + \cos x)}$ $= \frac{\sin^2 x + 1 + 2\cos x + \cos^2 x}{\sin x(1 + \cos x)}$ $= \frac{2 + 2\cos x}{\sin x(1 + \cos x)}$ $= \frac{2(1 + \cos x)}{\sin x(1 + \cos x)}$ $= \frac{2}{\sin x} \text{ (shown)}$ OR LHS $= \frac{\tan^2 x + (1 + \sec x)^2}{\tan x(1 + \sec x)}$ $= \frac{\tan^2 x + 1 + 2\sec x + \sec^2 x}{\tan x(1 + \sec x)}$ $= \frac{2\sec^2 x + 2\sec x}{\tan x(1 + \sec x)}$ $= \frac{2\sec x(\sec x + 1)}{\tan x(1 + \sec x)}$ $= \frac{2}{\cos x} \div \frac{\sin x}{\cos x}$ $= \frac{2}{\sin x} \text{ (shown)}$	M1: $\tan x = \frac{\sin x}{\cos x}$ & $\sec x = \frac{1}{\cos x}$ M1: simplify both [] correctly M1: add the fractions and expand correctly M1: factorise numerator AG1 OR M1: add fractions correctly M1: expand and add correctly M1: factorise numerator M1: $\sec x = \frac{1}{\cos x}$ & $\tan x = \frac{\sin x}{\cos x}$ AG1	AO2
-----------------	---	--	------------

10 ii	$\frac{\tan x}{1 + \sec x} + \frac{1 + \sec x}{\tan x} = 1 + 3\sin x$ $\frac{2}{\sin x} = 1 + 3\sin x$ $3\sin^2 x + \sin x - 2 = 0$ $(3\sin x - 2)(\sin x + 1) = 0$	M1: form quadratic eqn M1: factorisation or general formula	AO1
------------------	---	---	------------

$\sin x = \frac{2}{3}$ or $\sin x = -1$ $x = 41.8^\circ, 138.2^\circ$ or $x = 270^\circ$ (reject (to 1 dp) as $\tan 270^\circ$ is undefined)	A1, A1	
--	--------	--

11 i	$y = (9-x)(x-3)$ Sub $(k, k-3)$ into $y = (9-x)(x-3)$ $k-3 = (9-k)(k-3)$ * $k-3 = 9k-27-k^2+3k$ $k^2-11k+24=0$ $(k-3)(k-8)=0$ $k=3(N.A.)$ or $k=8$ *OR $(9-k)(k-3)-(k-3)=0$ $(k-3)(9-k-1)=0$ $(k-3)(8-k)=0$ $k=3(N.A.)$ or $k=8$	M1 substitution M1 form quadratic eqn M1 factorisation AG1 must state N.A. for $x=3$	AO3
-----------------------	--	---	-----

11 i	Let $y = 0$ $(9-x)(x-3) = 0$ $x = 3$ or $x = 9$ x-coordinate of $B = 9$ x-coordinate of $A = \frac{3+9}{2} = 6$ ** OR use $\frac{dy}{dx} = -2x + 12$ At turning point, $\frac{dy}{dx} = 0$ $-2x + 12 = 0$ $x_A = 6$  Area of $APCQ = 5 \times 2 = 10 \text{ units}^2$ Area $CQB = \int_8^9 (-x^2 + 12x - 27) dx$ $= \left[-\frac{1}{3}x^3 + 6x^2 - 27x \right]_8^9$ $= \left[-\frac{1}{3}(9^3) + 6(9^2) - 27(9) \right] - \left[-\frac{1}{3}(8^3) + 6(8^2) - 27(8) \right]$ $= 2\frac{2}{3} \text{ units}^2$ Shaded area $= 10 + 2\frac{2}{3} = 12\frac{2}{3} \text{ units}^2$	M1 : either x_A or x_B B1 (area of rectangle) M1: Integrate all terms correctly A1 A1	AO2
12 i	Gradient of $PQ = \text{gradient of } QR = \frac{1}{2}$ Eqn of PQ: $y - 3 = \frac{1}{2}(x + 4)$ $y = \frac{1}{2}x + 5$ ----- (1) Gradient of $QR = -2$ Eqn of QR: $y - 2 = -2(x - 4)$ $y = -2x + 10$ (1)=(2):	B1 M1: $m_1 m_2 = -1$ M1 (Equation of QR)	AO1

	$\frac{1}{2}x + 5 = -2x + 10$ $x = 2$ $\therefore Q(2, 6)$	M1: form simultaneous eqns A1	
12 ii	In eqn (1), Let $x=0$, $y=5$ $\therefore OT = 5$ units $RT = \sqrt{(4-0)^2 + (2-5)^2}$ $RT = \sqrt{25} = 5$ Since $OT = RT = 5$ units, $\triangle ORT$ is isosceles.	B1 B1 AG1	AO3
12 iii	Let $S = (a, b)$ By inspection: $S = (0-8, 0+1) = (-8, 1)$ OR Midpoint of $RS =$ Midpoint of OP $\left(\frac{a+4}{2}, \frac{b+2}{2}\right) = \left(-\frac{4}{2}, \frac{3}{2}\right)$ $a+4 = -4$ & $b+2 = 3$ $a = -8$ $b = 1$ Hence coordinates of $S = (-8, 1)$.	B1	AO2
12 iv	Area of trapezium OPQR $= \frac{1}{2} \begin{vmatrix} 0 & -4 & 2 & 4 & 0 \\ 0 & 3 & 6 & 2 & 0 \end{vmatrix}$ $= \frac{1}{2} -24 + 4 - 24 - 6 $ $= \frac{1}{2} -50 $ $= 25 \text{ units}^2$	M1 A1	AO1



Looking for a Home Tutor? WhatsApp us now at: +65 8820 0100
www.singaporehometutors.com